

# SSCM 1023 MATHEMATICAL METHODS I

## TOPIC: IMPROPER INTEGRAL

SHAZIRAWATI MOHD PUZI

ZUHAILA ISMAIL

&

NORZIEHA MUSTAPHA

DEPARTMENT OF MATHEMATICAL SCIENCES, UTM  
JB

# IMPROPER INTEGRALS

## 5.1 L'Hopital Rule

5.1.1 L'Hopital Rule for  $0/0$

5.1.2 L'Hopital Rule for  $\infty/\infty$

## 5.2 Improper Integrals

5.2.1 Improper Integral Type 1

5.2.2 Improper Integral Type 2

## 5.3 References

## 5.1 L'Hopital Rule

If you are doing any limit and you get something in the form of  $0/0$  or  $\infty/\infty$ , then you should probably try to use L'Hopital rule. The basic idea of L'Hospital rule is simple.

Consider the limit

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)}.$$

If both the numerator and the denominator are finite at  $a$  and  $g(a) \neq 0$ , then

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{f(a)}{g(a)}.$$

**Example 1:**

$$\lim_{x \rightarrow 3} \frac{x^2 + 1}{x + 2} = \frac{10}{5} = 2.$$

But what happen if both the numerator and the denominator tend to zero?

It is not clear what the limit is. In fact, depending on what functions  $f(x)$  and  $g(x)$  are, the limit can be anything at all!

### 5.1.1 L'Hopital Rule for 0/0

Suppose  $\lim f(x) = \lim g(x) = 0$ . Then

1. If  $\lim \frac{f'(x)}{g'(x)} = L$ , then

$$\lim \frac{f(x)}{g(x)} = \lim \frac{f'(x)}{g'(x)} = L.$$

2. If  $\lim \frac{f'(x)}{g'(x)}$  tends to  $+\infty$  or  $-\infty$  in the limit, then

so does  $\lim \frac{f(x)}{g(x)}$ .

***Example 2:***

Find  $\lim_{x \rightarrow 0} \frac{\sin x}{x}$  by L'Hopital rule.

***Example 3:***

Find  $\lim_{x \rightarrow 1} \frac{2 \ln x}{x - 1}$ .

***Example 4:***

Find  $\lim_{x \rightarrow 0} \frac{e^x - 1}{x}$ .

***Example 5:***

Find  $\lim_{x \rightarrow 0} \frac{1 - \cos x}{x}$ .

Note: If the numerator and the denominator both tend to  $+\infty$  or  $-\infty$ , L'Hopital rule still applies.

### 5.1.2 L'Hopital Rule for $\infty/\infty$

Suppose  $\lim f(x)$  and  $\lim g(x)$  are both infinite. Then

1. If  $\lim \frac{f'(x)}{g'(x)} = L,$

then  $\lim \frac{f(x)}{g(x)} = \lim \frac{f'(x)}{g'(x)} = L.$

2. If  $\lim \frac{f'(x)}{g'(x)}$  tends to  $+\infty$  or  $-\infty$  in the limit, then

so does  $\lim \frac{f(x)}{g(x)}.$

**Example 6:**

Find  $\lim_{x \rightarrow \infty} \frac{x^2}{e^x}.$

**Example 7:**

Find  $\lim_{x \rightarrow \infty} \frac{(\ln x)^2}{x}.$

## 5.2 Improper Integrals

The definite integral

$$\int_a^b f(x) dx$$

is known as *improper integral* if either

- 1) one or both limits are infinite, or
- 2)  $f(x)$  is undefined at certain points on/in the interval.

Note: We called case:

- 1) as Type I
- 2) as Type II

## 5.2.1 Improper Integral Type I

1) If  $f(x)$  is continuous in the interval  $[a, \infty)$ ,

$$\text{then } \int_a^{\infty} f(x) dx = \lim_{T \rightarrow \infty} \int_a^T f(x) dx.$$

2) If  $f(x)$  is continuous in the interval  $(-\infty, b]$ ,

$$\text{then } \int_{-\infty}^b f(x) dx = \lim_{T \rightarrow -\infty} \int_T^b f(x) dx.$$

Note: the improper integrals in 1) and 2) is said to *converge* if the limit exists and *diverge* if the limit does not exist.

3) If  $f(x)$  is continuous in the interval  $(-\infty, \infty)$ ,

$$\text{then } \int_{-\infty}^{\infty} f(x) dx = \int_{-\infty}^c f(x) dx + \int_c^{\infty} f(x) dx$$

with any real number  $c$ .

Note: the improper integrals in 3) is said to *converge* if both terms converge and *diverge* if either term diverges.



***Example 8:***

Determine whether the following integrals are convergent or divergent:

1) 
$$\int_0^{\infty} e^{-2x} dx$$

2) 
$$\int_0^{\infty} xe^{-x} dx$$

3) 
$$\int_{-\infty}^2 \frac{dx}{5-2x}$$

4) 
$$\int_{-\infty}^{\infty} \frac{x}{1+x^2} dx$$

## 5.2.2 Improper Integral Type II

- 1) If  $f(x)$  is continuous on  $[a, b)$ , and discontinuous at  $b$ , then

$$\int_a^b f(x) dx = \lim_{T \rightarrow b^-} \int_a^T f(x) dx.$$

- 2) If  $f(x)$  is continuous on  $(a, b]$ , and discontinuous at  $a$ , then

$$\int_a^b f(x) dx = \lim_{T \rightarrow a^+} \int_T^b f(x) dx.$$

Note: the improper integrals in 1) and 2) is said to *converge* if the limit exists and *diverge* if the limit does not exist.

- 3) If  $f(x)$  has discontinuity at  $c$ , where  $a < c < b$ , then

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx.$$

Note: the improper integrals in 3) is said to *converge* if both terms converge and *diverge* if either term diverges.

***Example 9:***

Determine whether  $\int_0^1 \frac{x}{\sqrt{1-x^2}} dx$  converge or diverge.

***Example 10:***

Determine whether  $\int_1^2 \frac{dx}{1-x^2}$  converge or diverge.

***Example 11:***

Find  $\int_{-1}^1 \ln x dx$  if possible.

## 5.3 References

1. George B. Thomas, Maurice D. Weir, Joel R. Hass, and Frank R. Giordano. 2005. *Thomas' Calculus Early Transcendental (11th Edition) (Thomas Series)*. Addison-Wesley Longman Publishing Co., Inc., Boston, MA, USA.
2. Abdul Wahid Md Raji, Hamisan Rahmat, Ismail Kamis, Mohd Nor Mohamad, Ong Chee Tiong. *Engineering mathematics I*, Penerbit UTM, 2012.